

## **Buy Finasteride 2.5mg Tablets Online from The Hair Dr**

### **Product Information**

Finasteride Accord is a proven treatment option for male pattern baldness, boasting a comparable efficacy profile to Propecia, with the added benefit of superior value. As a branded medication containing finasteride as its active ingredient, Finasteride Accord offers a reliable and effective solution for individuals seeking to address hair loss.

Finasteride has undergone extensive clinical research and investigation, solidifying its position as a leading treatment option for male pattern baldness. Studies have consistently demonstrated that finasteride effectively halts hair loss in a remarkable 90% of men, with the majority of individuals experiencing a significant reduction in hair thinning.

This treatment is specifically intended for men only.

Finasteride 1mg tablets have been rigorously tested and approved for use in the UK for the treatment of male pattern baldness.

When taken daily for a minimum of 3-6 months, this medication has been shown to improve the visible appearance and thickness of hair. However, it is essential to note that individual results may vary depending on age and the extent of existing hair loss.

It is crucial to adhere to the recommended dosage instructions: taking only ONE tablet daily. Exceeding this dose will not enhance the efficacy of Finasteride 1mg tablets and may potentially lead to adverse effects. Please consult the label instructions carefully and follow the recommended regimen.

### **Directions for Use**

For optimal results, patients taking Finasteride 1mg tablets should take one tablet once daily, with or without food. It typically takes a period of three to six months, with daily administration, before noticeable stabilization of hair loss can be expected to occur.

### **Side Effects**

Like all medications, Finasteride 1mg Accord tablets carry a risk of side effects. While some may be more common than others, most typically subside with continued use. If side effects persist, it's essential to report them to us at The Hair Dr, your pharmacist or your doctor.

It is crucial to seek immediate medical attention if you experience any of the following severe side effects:

- Swelling of the face, lips, tongue, or throat
- Difficulty swallowing
- Rash on the body or face, accompanied by itching and hives
- Breathing difficulties

In the event of such an occurrence, discontinue treatment with finasteride tablets and consult with your doctor or visit the nearest emergency department as soon as possible.

It is essential to promptly notify your doctor if you experience any of the following symptoms:

- Lumps, pain, or enlargement of breast tissue or nipple discharge. These signs may indicate a serious condition, including breast cancer, which can affect men as well as women.

**Package Leaflet: Information for the user Finasteride 1 mg Film-coated Tablet**

**Finasteride**

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

**What is in this leaflet?**

1. What Finasteride 1 mg Tablets are and what they are used for
2. What you need to know before you take Finasteride 1 mg Tablets
3. How to take Finasteride 1 mg Tablets
4. Possible side effects
5. How to store Finasteride 1 mg Tablets
6. Contents of the pack and other information

**1) What is Finasteride and what is it used for?**

Finasteride 1 mg Tablets contain finasteride. Finasteride belongs to the group of medicines called 5- alpha reductase inhibitors.

Your doctor has prescribed Finasteride 1 mg Tablets to address your male pattern hair loss, also referred to as androgenetic alopecia. As directed, these tablets have been designed to prevent further hair loss in men. Specifically, Finasteride 1 mg Tablets are suitable for individuals experiencing mild to moderate hair loss, but not complete baldness. By selectively inhibiting the enzyme Type II 5 $\alpha$ -reductase, which plays a crucial role in regulating the hair follicle, these tablets aim to promote a healthier and more sustainable hair growth pattern.

Finasteride 1 mg Tablets exert their therapeutic effects by selectively reducing dihydrotestosterone (DHT) levels in the scalp, a key contributor to male pattern hair loss. By mitigating DHT's influence, these tablets aim to arrest and potentially reverse the balding process, thereby preventing further hair loss and promoting a healthier hair growth profile.

**2) What is Finasteride and what is it used for?**

---

### **Do not take Finasteride 1mg tablets:**

- If you are allergic to Finasteride or any of the other ingredients of this medicine
- If you are a Child / Adolescent
- If you are Female
- Individuals currently taking finasteride or any other 5 $\alpha$ -reductase inhibitor for benign prostatic hyperplasia (BPH) or other indications should not use Finasteride 1 mg Tablets, as the concomitant use of these medications may lead to potential interactions and adverse effects.

### **Warnings and Precautions**

Talk to your doctor or pharmacist before starting a course of Finasteride.

Finasteride may impact male fertility. Patients contemplating fathering a child are advised to consider discontinuing treatment with Finasteride 1 mg Tablets, as the medication's effects on fertility have not been fully established and may lead to potential reproductive harm.

Prior to initiating therapy with Finasteride 1 mg Tablets, a baseline serum prostate-specific antigen (PSA) measurement should be obtained, and regular PSA assessments should be performed during treatment to monitor for potential changes in PSA levels.

### **Mood Alterations and Depression**

The medication Finasteride 1 mg Tablets has been linked to mood alterations, including depressed mood, depression, and in rare cases, suicidal thoughts. If you experience any of these symptoms, it is essential to discontinue use and consult with your doctor as soon as possible to receive further medical guidance.

### **Other medications and Finasteride**

It is crucial to inform your doctor or pharmacist about any medications you are currently taking, have recently taken, or may take in the future. While Finasteride 1 mg Tablets typically do not interact with other medications, there is limited

information on their use in combination with topical minoxidil for male pattern hair loss, and its use is not recommended.

### **Pregnancy, breast-feeding and fertility**

Finasteride 1 mg Tablets are specifically designed for men only.

Women, especially those who are or may be pregnant, should not handle the tablets, especially if broken or crushed, as there is a risk of the medication being absorbed through the skin or ingested, which could lead to birth defects in male foetuses. Additionally, if a patient's partner is or may become pregnant, they should either avoid exposure to semen or discontinue treatment with finasteride to prevent any potential harm.

### **Important information about some of the ingredients of Finasteride 1 mg Tablets:**

The Finasteride 1 mg Tablets contain lactose, and individuals with lactose intolerance should consult with their doctor before taking the medication.

### **3) How to take Finasteride 1mg Tablets**

The medication should be taken exactly as directed by your doctor, and if you're unsure, consult with your doctor or pharmacist.

The recommended dose is one 1mg tablet daily, which can be taken with or without food.

The tablets should be swallowed whole and not divided or crushed. It's important to note that taking more than one tablet a day will not enhance its effectiveness.

Additionally, male pattern hair loss is a gradual process, and it may take 3-6 months to notice results. Your doctor will help monitor the treatment's effectiveness, and it's crucial to continue taking the medication as prescribed.

### **Patients with impaired liver function**

The safety and efficacy of Finasteride 1 mg Tablets in patients with impaired liver function have not been established through clinical trials or experience.

---

## **Patients with impaired kidney function**

No dosage adjustment is needed for Finasteride 1 mg Tablets in patients with impaired kidney function.

### **If you take more Finasteride 1 mg Tablets than you should:**

If you take more Finasteride 1 mg Tablets than prescribed or if someone else accidentally takes some, seek immediate medical attention at the nearest emergency room. Bring any remaining tablets or empty packaging for easier identification.

### **If you forget to take Finasteride 1 mg Tablets:**

If you miss a dose of Finasteride 1 mg Tablets, simply take the next scheduled dose as usual. Do not take a double dose to make up for the missed dose.

### **If you stop taking Finasteride 1 mg Tablets:**

To achieve maximum benefit, it is recommended to continue taking Finasteride 1mg Tablets as directed. Stopping treatment may result in loss of previously gained hair within 9-12 months.

If you have any questions or concerns, consult your doctor or pharmacist for guidance.

<h2><b>4) How to take Finasteride 1mg Tablets</b></h2>
--

As with any medication, Finasteride 1 mg Tablets can cause side effects, although not everyone may experience them. Some side effects may be temporary and may resolve with continued treatment or disappear when treatment is stopped.

If you experience any of the serious side effects, stop taking Finasteride 1 mg Tablets immediately and seek medical attention: symptoms of an allergic reaction, such as swelling of the face, tongue, and throat, difficulty swallowing, hives, and breathing difficulties; or depression, characterized by severe sadness and feelings of unworthiness.

If you experience any changes in your breast tissue, including lumps, pain, enlargement of the breast tissue, or nipple discharge, report these changes to your

doctor immediately, as they may be signs of a serious condition such as breast cancer.

#### Uncommon side effects (affecting 1 in 100)

- Reduced libido / desire to have sex
- Difficulty achieving an erection, problems with ejaculation such as a decrease in semen release during sex
- Depression

#### **5) How to store Finasteride 1mg Tablets**

Keep Finasteride 1 mg Tablets out of the reach and sight of children. The medicine does not require special storage conditions and should not be used after the expiry date, which is indicated on the carton as a month and day. Additionally, dispose of unused medicine through a responsible means, such as by asking your pharmacist for guidance, to protect the environment.

#### **6) Contents of the pack and other valuable information**

The active ingredient in Finasteride 1 mg Tablets is finasteride, which is present in a dosage of 1mg per tablet. The tablet core contains a combination of excipients, including lactose, cellulose, starch, sodium starch glycolate, lauryl macroglycerides, and magnesium stearate. The film coating consists of hypromellose, titanium dioxide, macrogol 6000, iron oxide red, and iron oxide yellow.